Check List the journal of biodiversity data
NOTES ON GEOGRAPHIC DISTRIBUTION

The Surinam Cockroach, *Pycnoscelus surinamensis* (Linnaeus, 1758) (Insecta, Blattodea, Blaberidae) in Mexico: new records with the help of citizen science

Manuel de Luna[1,2,3], Julio C. Estrada-Álvarez[1,4], Reinier Núñez-Bazán[1,5], Roberto García-Barrios[6], Ismael A. Hinojosa Díaz[2], Gerardo Cuéllar-Rodríguez[2]

1 *Entomological Research, Metepec, Estado México, México*
2 *Instituto de Biología, Universidad Nacional Autónoma de México, Ciudad de México, México*
3 *Facultad de Ciencias Forestales, Universidad Autónoma de Nuevo León, Nuevo León, México*
4 *Museo Universitario de Historia Natural "Dr. Manuel M. Villada", Universidad Autónoma del Estado de México, México*
5 *Centro de Investigación en Biodiversidad y Conservación, Universidad Autónoma del Estado de Morelos, Morelos, México*
6 *Facultad de Ciencias Biológicas, Universidad Autónoma de Nuevo León, Nuevo León, México*

Corresponding author: *Reinier Núñez-Bazán (rnbazan7@gmail.com)*

Abstract. *Pycnoscelus surinamensis* (Linnaeus, 1758), Surinam Cockroach, is considered introduced to the Western Hemisphere, where it is a species of commercial and veterinary importance. In Mexico, *P. surinamensis* has been recorded from 17 states, and we newly report it in the states of Campeche, Michoacán, Puebla, and Querétaro, as well as the Ciudad de México, based on the examination of 184 specimens deposited in scientific collections. We also report it from the states of Aguascalientes, Baja California, Chihuahua, Estado de México, Guanajuato, Hidalgo, Sonora, and Tabasco based on our review of 952 photographic records from the citizen-science platform iNaturalist.

Key words. Blattodean, cosmopolitan species, introduced species, parasite, pest species, Pycnoscelinae, veterinary importance

de Luna M, Estrada-Álvarez JC, Núñez-Bazán R, García-Barrios R, Hinojosa-Díaz IA, Cuéllar-Rodríguez G (2025) The Surinam Cockroach *Pycnoscelus surinamensis* (Linnaeus, 1758) (Insecta, Blattodea, Blaberidae) in Mexico: new records with the help of citizen science. Check List 21 (2): 406–412. https://doi.org/10.15560/21.2.406

INTRODUCTION

The family Blaberidae comprises around 1,300 species of live-bearing cockroaches that usually have compact bodies and very short cerci which may sometimes appear absent. This family is very variable in its overall morphology, size, and lifestyle (Djernæs 2018). It is represented in continental North America by 16 genera (Atkinson et al. 1991; Estrada-Álvarez 2013; Paiero and Marshall 2014). Particularly, the genus *Pycnoscelus* Scudder, 1862, which is native of Southeast Asia, is present in all the countries of North America (Canada, Mexico, and the USA). One of its species, *Pycnoscelus surinamensis* (Linnaeus, 1758), Surinam Cockroach (Figure 1A, B), is a synanthropic invader and is considered cosmopolitan. It is an excellent colonizer that readily becomes established in new locations via the introduction of a single female (Roth 1998; Bell et al. 2007).

In Mexico, *P. surinamensis* has been recorded from 17 states: Baja California Sur (Hebard 1921; Palacios and Jiménez 1997); Chiapas (Fisk 1977; Reddell 1981; Palacios-Vargas et al. 2015); Coahuila (Cueto-Medina et al. 2015; Hernández-Rodríguez et al. 2013a, 2013b, 2014, 2015, 2016); Colima (Roth 1998); Durango (Arevalo-Arriaga 2012); Guerrero (Roth 1998); Jalisco (Hebard 1921; Cupul-Magaña et al. 2016); Morelos (Hoffman et al. 1986; Estrada-Álvarez and Guadarrama 2013; Núñez-Bazán et al. 2021); Nayarit (Ramírez-Olvera 1999); Nuevo León (Santoyo-de-Estéfano et al. 2014); Oaxaca (Hebard 1932); Quintana Roo (Lucio-Palacio and Noh-Gómez 2023); San Luis Potosí (Hebard 1932; Reddell 1981); Sinaloa (Hebard 1922); Tamaulipas (Reddell and Mitchell 1971); Veracruz (Saussure 1864; Walker 1868; Rehn 1902; Caudell 1914; Hebard 1921; Roth 1998); and Yucatan (Rehn 1902).

Pycnoscelus surinamensis is thought to have originated in the Indo-Pacific region, although its specific epithet implies another origin, Suriname. Populations of this species are made up almost entirely of thelytokous parthenogenetic females, although infertile males are produced, albeit extremely rarely (Roth 1967). Bisexual populations originally thought to belong to this taxon were elevated to species status as *Pycnoscelus indicus* (Fabricius, 1775), the sister lineage from which *P. surinamensis* arose (Roth 1967; Wang et al. 2025).

OPEN ACCESS

Academic editor: Leonardo de Oliveira Cardoso da Silva
Received: 26 December 2024
Accepted: 14 March 2025
Published: 4 April 2025

Copyright © The authors. This is an open-access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0)

This species is considered economically important as it can become a pest in greenhouses and plantations (Schwabe 1949; Roth and Willis 1960; Bell et al. 2007). Roth and Willis (1960) reported damage to several species of plants, and in some cases, it was as extensive as 30,000 to 300,000 individual plants destroyed. *Pycnoscelus surinamensis* is of veterinary importance as, alongside *P. indicus*, these two species are the only known intermediate hosts of the nematode *Oxyspiura mansoni* (Cobbold, 1879) (Spirurida, Thelaziidae), a parasite whose adults reside in the eyes of various birds, including domestic chickens (Schwabe 1949; Rodríguez-Tovar et al. 2008; Santoyo-de-Estéfano et al. 2014).

Here, we newly report *P. surinamensis* from several states in Mexico based on our examination of 184 specimens in entomological collections, as well as 952 research-grade observations from the citizen science platform iNaturalist. These data expand the knowledge of the distribution of this introduced insect in the country.

METHODS

Revision of specimens in collections. During a visit in June 2023 to the Colección Nacional de Insectos (CNIN) of the Universidad Nacional Autónoma de México, cockroach specimens deposited there were examined, and 51 specimens of *P. surinamensis* were found. Additionally, 42 specimens of *P. surinamensis* from the Laboratorio de Entomología of the Facultad de Ciencias Forestales (FCF) of the Universidad Autónoma de Nuevo León (UANL), 18 from the Colección Nacional de Insectos "Dr. Alfredo Barrera Marín, Museo de Historia Natural de la Ciudad de México, Chapultepec, CDMX, México (CNIABM), 15 from the Entomological Collection of Entomological Research (ER) and 58 from the American Museum of Natural History (AMNH) (Johnson et al. 2023) were also consulted.

Observations in iNaturalist. Mexican records in iNaturalist (2025) were reviewed on 28 February 2025. Blattodea records in general were checked. From those, observations of *P. surinamensis* were selected and identifications manually corroborated by examining the morphology and color patterns. Although *P. surinamensis* is somewhat variable in its coloration, both adults and nymphs are easily recognized from other North American cockroaches, even only from photographs. This makes *P. surinamensis* an ideal candidate to be monitored through citizen-science applications such as iNaturalist (see Identification below). Each of the confirmed iNaturalist records were marked as "research-grade" observations in the platform. Both adults and juveniles were included in our study. The map showing the distribution of the species in Mexico was made using SimpleMappr (Shorthouse 2010).

RESULTS

Order Blattodea Brunner von Wattenwyl, 1882
Family Blaberidae Saussure, 1865
Genus *Pycnoscelus* Scudder, 1862

***Pycnoscelus surinamensis* (Linnaeus, 1758)**
Figures 1, 2

Material examined. MEXICO — **Baja California Sur** • Lower California between San Jose del Cabo and Triunfo, Albatross Expedition; 1911; 5 ♀ and 19 juv., AMNH — **Campeche** • Champotón, El Zapote; 19.VI.1989; A. Cárdenas, L. Cervántes leg.; 1 ♀, UNAM-CNIN, PycSu001 • Bolano H.O.N. de Rejón; 21.VI.1986; F. Arias leg.; 1 ♀, UNAM-CNIN, PycSu002 — **Chiapas** • Huixtla; 8.VIII.1941; 1 juv., UNAM-CNIN, PycSu002 • Acapetahua; 17.IV1980; B. Melendes leg.; 1 ♀, UNAM-CNIN, PycSu003 • Río Monte Tapachula; 18.IX.1985; M. Vertiz leg.; 1 juv., UNAM-CNIN, PycSu004 — **Ciudad de México** • 5.X.1978; E. Marino leg.; 1 juv., UNAM-CNIN, PycSu005 • Del. Benito Juarez; 12.VIII.2006; A. Ibañez leg.; 1 juv., CNIABM 363-15 • 8.V.1968; A. Barrera leg.; 1 juv., CNIABM 363-16 — **Colima** • Isla Clarión; VII-XII.1987; E. Barrera leg.; 10 ♀, UNAM-CNIN, PycSu006–PycSu016 — **Durango** • San Juan de Guadalupe, San Juan de Guadalupe; 27.IX.2022; R. Nuñez-Bazán; 4♀, ER Pyc001-Pyc004 — Guerrero • Cocula, Guerrero; 17.II.1968; Q.B.P leg.; 1 juv., CNIABM 363-17 • Omiltemi, 1400 m a.s.l.; 16.V.1971; M. Cortés leg.; 1 juv., CNIABM 365-8 — **Jalisco** • Estación Biológica Chamela, 19°29'55.18"N, 105°02'39.26"W, 99 m a.s.l., 5–7.IX.2009; A. Zaldívar and H. Clebsch leg.; 1 ♀, UNAM-CNIN PycSu017 • Chamela; 5.XII.1976; Harry Brailovsky leg.; 1 ♀, UNAM-CNIN PycSu018 • Estación Biológica Chamela, La Huerta, Escolar Santa Cruz; 19°35'54.54"N, 105°02'55.43"W; 31.V.2022; A. Zaldívar and P. Benítez leg.; 1 ♀, UNAM-CNIN PycSu019 • 1 mile SW of "La Resolana"; 20.X.1950; Ray F. Smith leg.; 2 juv, AMNH • Ajijic; 20.VI.1968; G. y V. Halffter y P. Reyes leg.; 2 ♀, CNIABM 363-21, 363-22 — **Michoacán** • Morelia, Cerro Punhuato; 23.VIII.2014; López-López leg.; 1 ♀, UNAM-CNIN PycSu020 — **Morelos** • Xochitepec; 10.VII.1955; V. Guzmán leg.; 1 ♀, CNIABM 39160 • Teques-



Figure 1. *Pycnoscelus surinamensis* from Nuevo León, Mexico. **A.** Dorsal aspect of adult female with right tegmina and wing extended; abdomen shrunken due to being dried, making the wings and tegmina to appear longer. **B.** Dorsal aspect of subadult nymph. Scale bar: 5 mm.

quitenco; 5.IV.1969; Ma. T. Galván leg.; 1 ♀, CNIABM 365-7 • Loma Bonita, Cuernavaca; XI.2021; R. Núñez-Bazán leg.; 3 ♀, ER Pyc005-Pyc007 • Totolapan; 22.X.2023; J. Ramos leg.; 8 juv, ER Pyc008-Pyc015 — **Nayarit** • Isla María Magdalena, acueducto; 23.XI.1986; V. Melendez leg.; 1 juv., UNAM-CNIN PycSu021 • Río Santiago Colorado de la Mora, Arroyo Los Negros; 28.XI.1991; R. Barba and E. Barrera leg; 1 ♀, UNAM-CNIN PycSu022 • San Blas; 4–5.VIII.1947; B. Malkin leg.; 2 ♀, 4 juv, AMNH — **Nuevo León** • García; VII.1910; 1 ♀, UNAM-CNIN PycSu023 • Ciudad Universitaria, San Nicolás de los Garza; 13.I.2020, Roberto García-Barrios leg.; 2 ♀ and 5 juv., FCF-BLATT021 • Facultad de Ciencias Forestales UANL, Linares; 15.I.2021; Manuel de Luna leg.; 1 ♀, 4 juv., FCF-BLATT024 • Panteón Municipal, Santa Catarina; Manuel de Luna leg.; 1 ♀, FCF-BLATT028 • Cerro de la Silla (cascadas, Rincón de la Sierra), Guadalupe; 2018; Manuel de Luna leg.; 2 ♀, FCF-BLATT107 — **Oaxaca** • Huatulco; 28.IX–11.X.2021; A. Zaldívar and P. Benítez leg.; 1 ♀, UNAM-CNIN PycSu025 • Santa María Huatulco, predio FONATUR evento 3B, 15°43'51.2"N, 096°09'45.61"W; 31.VIII.2021; A. Zaldívar and P. Benítez leg.; 1 ♀, UNAM-CNIN PycSu026 • Huatulco, evento 1B, 15°43'38.68"N, 096°09'43.91"W; 21.VI–7.VII.2021; A. Zaldívar and P. Benítez leg.; 1 ♀, UNAM-CNIN PycSu027 • Tuxtepec; 11.I.1964; 1 ♀, UNAM-CNIN PycSu028 • Monteflor, Oaxaca; 1 ♀, UNAM-CNIN PycSu029 • Piedra Lumbre; 3.XI.1987; F. Arias leg.; 2 juv., UNAM-CNIN PycSu030, PycSu031 • San Geronimo; 5.II.1909; 1 ♀, AMNH • Tolosa; 23.VIII.1947; B. Malkin leg.; 1 ♀ and 2 juv., AMNH — **Puebla** • Villa Avila, Camacho; 24.I.1985; E. Marino leg.; 1 juv, UNAM-CNIN PycSu032 • Matamoros; 6.VII.1942; L. Vázquez leg.; 1 ♀, UNAM-CNIN PycSu033 • Cuetzalán; 26.X.2017; Jonathan Velazquez leg.; 1 ♀, UNAM-CNIN PycSu033 • Matamoros; 18.VII.1942; 1 ♀, UNAM-CNIN PycSu034 — **Querétaro** • Ciudad de Querétaro; 1–7.XI.2008; K.A. Brailovsky leg.; 1 ♀, UNAM-CNIN PycSu035 — **Quintana Roo** • Isla Mujeres, Quintana Roo; 16.V.1961; F. Aguilar leg.; 6 juv., CNIABM 39203-08 — **San Luis Potosi** • El Salto Sn. Luis Potosi, 400 m a.s.l.; 22.V.1944; C. Bolivar leg; 2♀, 1 juv., CNIABM PycSu036 • Tamazunchale; 17.VII.1946; J. & D. Pallister leg.; 2 ♀, AMNH — **Sinaloa** • Mazatlán; 16–31.XII.1946; J. A. Kusche leg.; 1 ♀, AMNH • Mazatlán; Hy. Edwards leg.; 1 ♀, AMNH — **Tamaulipas** • Cerro Bernal; 21.XII.1949; F. Biaggi leg.; 1 juv., UNAM-CNIN PycSu037 • Reserva de la Biósfera "El Cielo", Gomez Farías; 15.X.2020; Roberto García-Barrios leg.; 2 ♀ and 2 juv., FCF-BLATT011 • Reserva de la Biósfera "El Cielo", Gomez Farías; 15.VIII.2021; Manuel de Luna leg.; 1 ♀, FCF-BLATT017 • Reserva de la Biósfera "El Cielo", Gomez Farías; 15.VII.2021; Manuel de Luna leg.; 3 ♀, FCF-BLATT030 • Reserva de la Biósfera "El Cielo", Gomez Farías; 24.X.2020; Roberto García-Barrios leg., 1 ♀, FCF-BLATT077 • Reserva de la Biósfera "El Cielo", Gomez Farías; 15.VI.2021; Manuel de Luna leg.; 4 ♀, FCF-BLATT086 • Reserva de la Biósfera "El Cielo", Gomez Farías; 15.XI.2020; Manuel de Luna leg.; 1 ♀, FCF-BLATT106 • Reserva de la Biósfera "El Cielo", Gomez Farías; 15.XI.2020; Manuel de Luna leg.; 1 ♀, FCF-BLATT108 • Reserva de la Biósfera "El Cielo", Gomez Farías; 15.VI. 2021; Manuel de Luna leg.; 8 ♀, FCF-BLATT143 — **Veracruz** • Huasteca Veracruzana; VIII.1958; 2 ♀, UNAM-CNIN PycSu038-PycSu039 • Estación de Biología Tropical "Los Tuxtlas"; 15.X.1985; E. Ramírez leg.; 1 ♀, UNAM-CNIN PycSu040 • Plan del Río; 1 ♀, UNAM-CNIN PycSu041 • Xalapa; 22.X.1983; J. Peña leg.; 1 juv., UNAM-CNIN PycSu042 • El Vigía, Santiago Tuxtla; 20.IX.1963; 6 juv., UNAM-CNIN PycSu043 • Ocotal Texisapa, 8.XII.1985, R. Barba leg.; 2 juv., UNAM-CNIN PycSu044 • Estación Biológica Tropical "Los Tuxtlas", San Andrés Tuxtla, 25.VII.2022; Manuel de Luna leg.; 2 ♀ and 2 juv., FCF-BLATT054 • Jesus Carranza; VI.1944; M. Guerra leg.; 1 ♀, AMNH. • Dos Amates; 27.XI.1967; P. Reyes leg.; 2 juv., CNIABM 362-8, 362-9 — **Yucatán** • Uxman, 20°24'00.97"N, 089°47'53.4"W; 11.VI.1994; E. Barrera leg.; 1 ♀, UNAM-CNIN PycSu045 • Chichen Itza; 16.VII.1952 to 19.VIII.1952; J. & D. Pallister leg.; 7 ♀, AMNH • Chuminopolis; 7.VIII.1952; J. & D. Pallister leg.; 1 ♀, AMNH • Colonia Yucatán; 6.VIII.1952 and



Figure 2. Distribution of *Pycnoscelus surinamensis* in Mexico. Records of the examined specimens and the photographic records are represented with yellow triangles. Blue-filled states denote the presence of this species as stated in previous literature records (see Introduction).

17.VIII.1952; colls. J. and D. Pallister leg.; 2 ♀, AMNH • Itzimna; 3–7.IX.1964; J. C. & D. Pallister leg.; 2 ♀ and 1 juv., AMNH • Chumihopilis; 20.VII.1964 and 8.VIII.1965; J. C. & D. Pallister leg.; 3 ♀, AMNH • Piste; 3–5.VI.1959; P. & C. Vaurie leg.; 1 ♀, AMNH.

iNaturalist observations. There were found 952 confirmed observations of *P. surinamensis* from Mexico in the platform, including seven from Aguascalientes, three from Baja California, 21 from Chihuahua, five from Estado de México, 25 from Guanajuato, 41 from Hidalgo, 22 from Sonora, and six from Tabasco. These constitute additional new state records. No records were found in the states of Tlaxcala and Zacatecas. See Supplementary file.

Identification. We based our identifications on the works of Roth (1967, 1998), Helfer (1987), Anisyutkin (2002, 2018), Choate (1998), and Lucio-Palacio and Noh-Gómez (2023), as well as the examination of a large series of specimens in the CNIN-UNAM and the FCF-UANL collections. This is a medium-sized cockroach (up to 3.23 cm in total length) with discreet coloration. The pronotum is black, with the anterior margin having yellow or cream-colored tinges; these can take the form of a band and extend to the lateral margins (Lucio-Palacio and Noh-Gómez 2023), be reduced (Figure 1A), or even be absent in melanistic individuals. Tegmina are typically light brown or ochre, with the proximal portion of the costal region usually being lighter than the rest (Figure 1A); in melanistic individuals the tegmina are completely black. The hindwings are dull brown and subhyaline (Figure 1A). The tergites are mostly black and very shiny, and these can have a yellow anterior margin, as well as yellow lateral patterns (Figure 1A), but this is variable; in melanistic individuals the terguites can be completely black. Both sexes have the same coloration and overall morphology, but most often only females can be found, as infertile males are exceedingly rare. Males have an asymmetrical subgenital plate and tegmina and wings which noticeably surpass the abdominal apex. The combination of these characteristics set *Pycnoscelus* apart from the rest of the North American Blaberidae. Nymphs are black (reddish brown in old, dried specimens), concolorous, and ostensibly shiny, except for the last four tergites and the supra-anal plate, which are opaque and roughened (Figure 1B) and often accumulating debris. This last character sets the nymphs of *P. surinamensis* apart from those of *Nauphoeta* Burmeister, 1838 and *Panchlora* Burmeister, 1838, which have similar morphology.

DISCUSSION

Pycnoscelus surinamensis had been reported in Mexico from 17 states (see Introduction). With the new records, it is now also recorded from Aguascalientes, Baja California, Campeche, Chihuahua, Ciudad de México, Estado de México, Guanajuato, Hidalgo, Michoacán, Puebla, Querétaro, Sonora and Tabasco. The only Mexican states without records of this species are Tlaxcala and Zacatecas. As has been recently explored with other polyneopterans (Estrada-Álvarez et al. 2023; de Luna et al. 2023), the use of citizen-science records proves to be extremely useful in filling knowledge gaps in the distribution of *P. surinamensis* in Mexico, and these observations are very important information for supplementing museum specimens. In this study, the records of *P. surinamensis* from Aguascalientes, Baja California, Chihuahua, Estado de México, Guanajuato, Hidalgo, Sonora and Tabasco were obtained exclusively from iNaturalist.

Currently, *P. surinamensis* does not figure as an invasive species in Mexico (DOF 2016; Lucio-Palacio and Noh-Gómez 2023); however, this must certainly be changed, as it has been proven to be a vector of a parasite with commercial and ecological importance and can deal hefty damage to both ornamental plants and crops. It also spreads very easily by parthenogenetic reproduction. All these characteristics make it a serious threat wherever it is introduced.

ACKNOWLEDGEMENTS

We thank the Society of Systematic Biologists for providing MdL with funding through the Mini-ARTS grant awarded to the project "Preliminary research on four cockroach families (Blaberidae, Blattellidae, Nyctiboridae, and Pseudophyllodromiidae) from Mexico". We also want to thank Alejandro Zaldívar Riverón and Christina Mayorga Martínez for allowing the first and third author to revise the specimens at the CNIN-UNAM. MdL thanks CONAHCyT for the postdoctoral fellowship (CVU 1032171). Finally, our gratitude extends to the anonymous reviewers for the corrections and comments, which helped better this publication.

ADDITIONAL INFORMATION

Conflict of interest

The authors declare that no competing interests exist.

Ethical statement

No ethical statement is reported.

Funding

This study was financially supported by the Society of Systematic Biologists

Author contributions

Conceptualization: JCEA, MdL, RNB. Data curation: RGB. Formal analysis: MdL, RGB. Funding acquisition: JCEA, MdL, IAHD, GCR. Investigation: JCEA, MdL, RNB. Methodology: JCEA. Project administration: JCEA, MdL, RNB. Software: MdL. Resources: MdL, IAHD. Supervision: IAHD. Validation: MdL, JCEA, RNB. Writing – original draft: MdL. Writing – review & editing: MdL, JCEA, RNB, RGB, IAHD.

Author ORCID iDs

Julio C. Estrada Álvarez https://orcid.org/0000-0002-3097-0599
Manuel de Luna https://orcid.org/0000-0002-0746-0507
Reinier Núñez Bazán https://orcid.org/0000-0002-8622-3373
Roberto García Barrios https://orcid.org/0000-0001-6062-9001
Ismael A. Hinojosa Díaz https://orcid.org/0000-0002-5493-850X
Gerardo Cuéllar Rodríguez https://orcid.org/0000-0003-4969-611X

Data availability

All data that support the findings of this study are available in the main text.

REFERENCES

Anisyutkin LN (2002) Notes on cockroaches of the subfamilies Pycnoscelinae and Diplopterinae from south-east Asia with description of three new species (Dictyoptera: Blaberidae). Zoosystematica Rossica 10: 351–359.

Anisyutkin LN (2018) Little known Epilamprinae (Dictyoptera: Blaberidae) from the collections of the Muséum d'histoire naturelle de Genève. Part 3. Revue Suisse de Zoologie 125: 1–16. https://doi.org/10.5281/zenodo.1196003

Arévalo-Arriaga RL (2012) Identificación de especies de cucarachas de importancia urbana en el área sureste de Gómez Palacio, Durango. Tesis presentada como requisito parcial para obtener el título de Ingeniero Agrónomo Parasitólogo, Universidad Autónoma Agraria "Antonio Narro", Torreón, Coahuila, México, 57 pp.

Atkinson TH, Koehler PG, Patterson RS (1991) Catalog and atlas of the cockroaches (Dictyoptera) of North America north of Mexico. Miscellaneous Publications of the Entomological Society of America 78: 1–85: https://doi.org/10.4182/wtam6452.78.11

Beccaloni GW (2023) https://cockroach.speciesfile.org/. Cockroach Species File. Accessed on: 2023-10-26.

Bell WJ, Roth LM, Nalepa CA (2007) Cockroaches: ecology, behavior, and natural history. Johns Hopkins University Press, Baltimore, USA, 247 pp.

Caudell AN (1914) Some bromeliadicolous Blattidae from Mexico and Central America. Insecutor Inscitiae Menstruus 2: 76–80.

Choate PM (1998) A dichotomous key for the identification of the cockroach fauna (Insecta: Blattaria) of Florida. Department of Entomology and Nematology, University of Florida, Gainesville, USA, 18 pp.

Cueto-Medina SM, Castillo-Martínez A, Hernández-Rodríguez S, Gallegos-Robles MÁ, Sánchez-Ramos FJ, Ortega-Morales AI (2015) Presencia de *Salmonella* spp. en cucaracha americana *Periplaneta americana* l. (Blattodea:Blattidae) en Torreón, Coahuila, México. Entomología Mexicana 2: 762–766

Cupul-Magaña FG, McCann F, Escobedo-Galván AH, Estrada-Álvarez JC (2016) *Aspidoscelis lineattisima* (Cope, 1878). Diet. Mesoamerican Herpetology (Nature Notes) 3: 482–483.

de Luna M, Núñez-Bazán R, Estrada-Álvarez JC, García-Barrios R (2023) New records of the striped earwig *Labidura riparia* (Pallas, 1773) (Insecta: Dermaptera: Labiduridae) in Mexico. Revista Chilena de Entomología 49: 727–734. https://doi.org/10.35249/rche.49.4.23.06

Djernaes SM (2018) Biodiversity of Blattodea—the cockroaches and termites. In: Footitt RG, Adler PH (Eds.) Insect biodiversity: science and society, volume 2. John Wiley & Sons, Hoboken, USA, 359–387.

DOF [Diario Oficial de la Federación] (2016) Acuerdo por el que se determina la Lista de las Especies Exóticas Invasoras para México. https://www.dof.gob.mx/nota_detalle.php?codigo=5464456&fecha=07/12/2016#gsc.tab=0. Accessed on: 2023-10-30.

Estrada-Álvarez JC (2013) Primera lista de las cucarachas (Insecta: Dictyoptera: Blattaria) de México. Boletín de la Sociedad Entomológica Aragonesa 53: 267–284.

Estrada-Álvarez JC, de Luna M, Núñez-Bazán R, Sormani CG (2023) New state records of the firefly-mimicking cockroach *Achroblatta luteola* (Blanchard, 1843) (Blattodea: Blaberidae: Panchlorinae) in Mexico. Revista Chilena de Entomología 49: 741–745. https://doi.org/10.35249/rche.49.4.23.08

Estrada-Álvarez JC, Guadarrama CAR (2013) Nuevos registros de cucarachas (Blattodea) para México. Dugesiana 20: 49–53. https://doi.org/10.32870/dugesiana.v20i1.4081

Fisk FW (1977) Subterranean fauna of Mexico. Part III. Further results of the Italian zoological missions to Mexico, sponsored by the National Academy of Lincei (1973 and 1975). Accademia Nazionale dei Lincei 171: 267–274.

Hebard M (1921) Mexican records of Blattidae (Orthoptera). Transactions of the American Entomological Society 47: 199–220.

Hebard M (1922) Dermaptera and Orthoptera from the State of Sinaloa, Mexico: part I. Dermaptera and non-saltatorial Orthoptera. Transactions of the American Entomological Society 48: 157–196.

Hebard M (1932) New species and records of Mexican Orthoptera. Transactions of the American Entomological Society 58: 201–371.

Helfer JR (1987) How to know the grasshoppers, crickets, cockroaches, and their allies. Dover Publications, New York, USA, 363 pp.

Hernández-Rodríguez S, López- Hernández J, Valdés-Perezgasga MT, Sánchez-Ramos FJ (2014) Cucarachas en el municipio de Matamoros, Coahuila, México. Entomología Mexicana 1: 925–930.

Hernández-Rodríguez S, Ortega-Morales AI, Valdés-Perezgasga MT, Sánchez-Ramos FJ, López-Hernández J, Santillán-Santana J (2013b) Nuevos registros de cucarachas urbanas en Torreón, Coahuila, México (Insecta: Blattodea). Acta Zoológica Mexicana (Nueva Serie) 29: 428–430.

Hernández-Rodríguez S, Valdés-Perezgasga MT, López-Hernández J, García-Espinoza F, Castillo-Martínez A, Cueto-Medina SM (2015) Cucarachas (Hexapoda: Blattodea) del municipio de san Pedro de las colonias, Coahuila, México. Entomología Mexicana 2: 695–700.

Hernández-Rodríguez S, Valdés-Perezgasga MT, López-Hernández J, García-Espinoza F, Hernández-Hernández V, Cueto-Medina SM (2016) Cucarachas (Insecta: Blattodea) del municipio de Francisco I. Madero, Coahuila, México. Entomología Mexicana 3: 748–753

Hernández-Rodríguez S, Valdés-Perezgasga MT, Sánchez-Ramos FJ, López- Hernández J., Ortega-Morales AI, Cueto-medina SM (2013a) Cucarachas en el municipio de Torreón, Coahuila, México. Entomología Mexicana 12: 1297–1302.

Hoffmann A, Palacios-Varga JG, Morales-Malacara JB (1986) Manual de Bioespeleología (con nuevas aportaciones de Morelos y Guerrero, México). Universidad Nacional Autónoma de México, Mexico City, México, 274 pp.

iNaturalist (2023) Records of *Pycnoscelus surinamensis* from Mexico. https://www.inaturalist.org/observations?place_id=6793&taxon_id=84635. Accessed on: 2023-10-30.

Johnson C, Ruiz AM, Tsutsumi K, Larsen E, Jones L, Salas R (2023) Digitization PEN: Digital data from the terrestrial Polyneoptera at American Museum of Natural History. Version 1.7. https://doi.org/10.15468/sg7c8c. Accessed via GBIF.org on: 2023-11-08.

Lucio-Palacio CR, Noh-Gómez JM (2023) Nuevos registros, distribución e historia natural de la cucaracha invasora *Pycnoscelus surinamensis* (Linnaeus, 1758) (Blattodea: Blaberidae: Pycnoscelinae) en Quintana Roo, México. Revista Chilena de Entomología 49: 861–869. https://doi.org/10.35249/rche.49.4.23.19

Núñez-Bazán R, Estrada-Álvarez JC, Sormani CGH, Osorio-Beristain M (2021) New data on Blaberoidea Saussure, 1864 (Insecta: Blattodea) from Morelos, Mexico, with a new species and a checklist of cockroaches for the state. Revista Chilena de Entomología 47: 689–708. https://doi.org/10.35249/rche.47.4.21.05

Paiero SM, Marshall SA (2014) New Canadian and Ontario orthopteroid records, and an updated checklist of the Orthoptera of Ontario. Journal of the Entomological Society of Ontario 145: 61–76.

Palacios C, Jimenez M (1997) Presencia de cucarachas en viviendas de La Paz, Baja California sur México. Southwestern Entomologist 22: 243–246.

Palacios-Vargas JG, Juberthie C, Reddell JR (2015) Encyclopædia Biospeologica. Vol. II, México. Unión Mexicana de Agrupaciones Espeleológica/Mundos Subterráneos 25–26: 1–101.

Ramírez-Olvera MA (1999) Lista preiliminar de la espeleofauna de la Isla Isabel Maria (Oaxaca, México). Mundos Subterraneos 10: 47–54.

Reddell JR (1981) A review of the cavernicole fauna of Mexico, Guatemala, and Belize. Bulletin Texas Memorial Museum, University of Texas at Austin 27: 1–327.

Reddell JR, Mitchell RW (1971) A checklist of the cave fauna of Mexico. II. Sierra de Guatemala. Tamaulipas. In: Reddell JR, Mitchell RW (Eds.) Studies on the cavernicole fauna of Mexico. Bulletin of the Association for Mexican Cave Studies, Austin, USA, 181–215.

Rehn JAG (1902) A contribution to the knowledge of the Orthoptera of Mexico and Central America. Transactions of the American Entomological Society 29: 1–34.

Rodríguez-Tovar LE, Casas-Martínez A, Ramírez-Romero R, Nevárez-Garza AM, Zárate-Ramos JJ (2008) First report of *Oxyspirura* sp. in a captive fulvous owl (*Strix fulvescens*) in Mexico. Journal of Parasitology 94: 1430–1431. https://doi.org/10.1645/ge-1599.1

Roth LM (1967) Sexual isolation in the parthenogenetic cockroach *Pycnoscelus surinamensis* and application of the name *Pycnoscelus indicus* to its bisexual relative (Dictyoptera: Blattaria: Blaberidae: Pycnoscelinae). Annals of the Entomological Society of America 60: 774–779. https://doi.org/10.1093/aesa/60.4.774

Roth LM (1998) The cockroach genus *Pycnoscelus* Scudder, with a description of *Pycnoscelus femapterus*, sp. nov. (Blattaria: Blaberidae: Pycnoscelinae). Oriental Insects 32: 93–130. https://doi.org/10.1080/00305316.1998.10433770

Roth LM, Willis ER (1960) The biotic associations of cockroaches. Smithsonian Miscellaneous Collections 141: 1–470.

Santoyo-de-Estéfano FA, Espinoza-Leija RR, Zárate-Ramos JJ, Hernández-Velasco X (2014) Identification of *Oxyspirura mansoni* (Spirurida: Thelaziidae) in a free-range hen (*Gallus gallus domesticus*) and its intermediate host, Surinam Cockroach (*Pycnoscelus surinamensis*) in Monterrey, Nuevo Leon, Mexico. Acta Zoologica Mexicana 30: 106–113.

Saussure H (1864) Orthoptères de l'America moyenne, Mémoires pour servir a L'Histoire naturelle du Mexique des Antilles et des États-Unis quinque memorie. Hachette Livre Bnf, Paris, France, 276 pp.

Schwabe CW (1949) Observations on the life history of *Pycnoscelus surinamensis* (Linn.), the intermediate host in the Chicken Eye-worm in Hawaii. Proceedings of the Hawaiian Entomological Society 13: 433–436.

Shorthouse DP (2010) SimpleMappr, an online tool to produce publication-quality point maps. https://www.simplemappr.net/. Accessed on: 2023-10-30.

Walker F (1868) Catalogue of the specimens of Blattariae in the collection of the British Museum. British Museum, London, United Kingdom, 239 pp. https://doi.org/10.5962/bhl.title.8495

Wang YS, Dong XJ, Wang ZQ, Che YL (2025) Taxonomic update of *Pycnoscelus* Scudder, 1862 (Blattodea, Blaberidae, Pycnoscelinae), with descriptions of two new species from China. ZooKeys 1231: 45–67. https://doi.org/10.3897/zookeys.1231.141287